




PET/GB04/2659



INVESTOR IN PEOPLE

The Patent Office  
Concept House  
Cardiff Road  
Newport  
South Wales  
NP10 8QQ



I, the undersigned, being an officer duly authorised in accordance with Section 74(1) and (4) of the Deregulation & Contracting Out Act 1994, to sign and issue certificates on behalf of the Comptroller-General, hereby certify that annexed hereto is a true copy of the documents as originally filed in connection with the patent application identified therein.

In accordance with the Patents (Companies Re-registration) Rules 1982, if a company named in this certificate and any accompanying documents has re-registered under the Companies Act 1980 with the same name as that with which it was registered immediately before re-registration save for the substitution as, or inclusion as, the last part of the name of the words "public limited company" or their equivalents in Welsh, references to the name of the company in this certificate and any accompanying documents shall be treated as references to the name with which it is so re-registered.

In accordance with the rules, the words "public limited company" may be replaced by p.l.c., plc, P.L.C. or PLC.

Re-registration under the Companies Act does not constitute a new legal entity but merely subjects the company to certain additional company law rules.



Signed *P. McHoney*  
Dated 3 August 2004

**BEST AVAILABLE COPY**

**PRIORITY  
DOCUMENT**  
SUBMITTED OR TRANSMITTED IN  
COMPLIANCE WITH RULE 17.1(a) OR (b)

# Request for grant of a patent

(See the notes on the back of this form. You can also get an explanatory leaflet from the Patent Office to help you fill in this form)



22SEP03 E838686-5 D02732  
P01/7700 0.00-0322003.5

The Patent Office

Cardiff Road  
Newport  
South Wales  
NP9 1RH

1. Your reference  
A1590
2. Patent application number  
(The Patent Office will fill in this part)  
19 SEP 2003 0322003.5
3. Full name, address and postcode of the or of each applicant (underline all surnames)  
  
08642183 001  
Patents ADP number (if you know it)  
  
If the applicant is a corporate body, give the country/state of its incorporation  
  
ABERDEEN UNIVERSITY  
RESEARCH AND INNOVATION  
UNIVERSITY OFFICE  
KINGS COLLEGE  
ABERDEEN AB24 3FX  
UK
4. Title of the invention  
SPONGE TOXINS
5. Name of your agent (if you have one)  
  
"Address for service" in the United Kingdom to which all correspondence should be sent (including the postcode)  
  
ABLETT & STEBBING  
CAPARO HOUSE  
101-103 BAKER STREET  
LONDON  
W1U 6FQ  
  
Patents ADP number (if you know it) 00006551001 ✓
6. If you are declaring priority from one or more earlier patent applications, give the country and the date of filing of the or of each of these earlier applications and (if you know it) the or each application number  

Country	Priority application number (if you know it)	Date of filing (day / month / year)
---------	--	-------------------------------------
7. If this application is divided or otherwise derived from an earlier UK application, give the number and the filing date of the earlier application  

Number of earlier application	Date of filing (day / month / year)
-------------------------------	-------------------------------------
8. Is a statement of inventorship and of right to grant of a patent required in support of this request? (Answer 'Yes' if:  
a) any applicant named in part 3 is not an inventor, or  
b) there is an inventor who is not named as an applicant, or  
c) any named applicant is a corporate body.  
See note (d)) YES

# Patents Form 1/77

9. Enter the number of sheets for any of the following items you are filing with this form. Do not count copies of the same document

Continuation sheets of this form	0
Description	48
Claim(s)	4 <i>DL</i>
Abstract	0
Drawing(s)	20 + 20

10. If you are also filing any of the following, state how many against each item.

Priority documents	0
Translations of priority documents	0
Statement of inventorship and right to grant of a patent ( <i>Patents Form 7/77</i> )	0
Request for preliminary examination and search ( <i>Patents Form 9/77</i> )	0
Request for substantive examination ( <i>Patents Form 10/77</i> )	0
Any other documents ( <i>please specify</i> )	0

11.

I/We request the grant of a patent on the basis of this application.

Signature

Date

*ABLETT & STEBBING*

19 September 2003

12. Name and daytime telephone number of person to contact in the United Kingdom

G K ABLETT / S J SUÈR (0207-935-7720)

## Warning

After an application for a patent has been filed, the Comptroller of the Patent Office will consider whether publication or communication of the invention should be prohibited or restricted under Section 22 of the Patents Act 1977. You will be informed if it is necessary to prohibit or restrict your invention in this way. Furthermore, if you live in the United Kingdom, Section 23 of the Patents Act 1977 stops you from applying for a patent abroad without first getting written permission from the Patent Office unless an application has been filed at least 6 weeks beforehand in the United Kingdom for a patent for the same invention and either no direction prohibiting publication or communication has been given, or any such direction has been revoked.

## Notes

- If you need help to fill in this form or you have any questions, please contact the Patent Office on 0645 500505.
- Write your answers in capital letters using black ink or you may type them.
- If there is not enough space for all the relevant details on any part of this form, please continue on a separate sheet of paper and write "see continuation sheet" in the relevant part(s). Any continuation sheet should be attached to this form.
- If you have answered 'Yes' Patents Form 7/77 will need to be filed.
- Once you have filled in the form you must remember to sign and date it.
- For details of the fee and ways to pay please contact the Patent Office.

## Sponge Toxins

The present invention relates to the use of sponge toxins, in particular polymeric 1,3-alkylpyridinium salts 5 (poly-APS), for the reversible formation of membrane pores and a method for producing such pores.

Over the last 500-700 million years sessile marine sponges have developed a variety of strategies to survive 10 changing environmental conditions, potential problems with dispersal and competition with other organisms including invasion of their structure by micro-organisms and the emergence of potential predators. In addition to being able to regenerate and take on a single cell existence, sponges 15 have an extensive armoury of chemical defences, which prevent over-predation, facilitate establishment of a sponge colony and control colonisation of the surfaces of sponges by other marine organisms. Of the many distinct chemical weapons produced by sponges a number are alkylpyridinium salts that 20 have interesting biological properties that may be exploited. Halitoxins (1,3-APS oligomers) were originally identified from sponges of the Haploscerid genera such as Haliclona, Amphimedon and Callyspongia. Diverse biological activities have been identified for different halitoxin preparations. 25 These include cytotoxicity, neurotoxicity and inhibition of action potentials, stimulation of transmitter release, inhibition of  $K^+$  conductances and anticholinesterase activity.

The chemical and biological properties of some natural 30 pore forming halitoxins in a fraction with a mean molecular weight of 5 kDa have been previously isolated and characterised (Scott et al (2000) J Membrane Biol 176 119-131). These compounds, isolated from the sponge *Callyspongia*

*ridleyi*, were stable in aqueous solution and it was found that this cocktail of natural halitoxins depolarised cultured sensory neurones by irreversibly forming pores permeable to cations. These irreversible pores result in irreversible damage to the cell wall which is lethal to the cell. When applied to artificial bilayers toxin-evoked channel-like events were obtained with unitary conductances between 145-2280pS. Additionally, the halitoxins could release  $Ca^{2+}$  from intracellular stores.

10

Single stranded polynucleotide RNA and DNA have been shown to transverse lipid bilayers treated with *Staphylococcus aureus*  $\alpha$ -toxin or  $\alpha$ -hemolysin, which create pores of 1 to 2.5nm diameter. These *Staphylococcus aureus*  $\alpha$ -toxin or  $\alpha$ -hemolysin produce irreversible pores. Double stranded DNA has been transferred across lipid bilayers and into intact cells using lipid-micelle-mediated transfection (e.g. lipofection), electroporation and microinjection but these approaches yield variable results and can be difficult to control often requiring specialist equipment or laborious pretreatments and optimisation.

20

It is therefore an object of the present invention to attempt to overcome the disadvantages of the prior art.

25

According to the present invention, there is provided use of a composition comprising a sponge toxin for the reversible formation of a membrane pore.

30 It has now been established that a pore can be reversibly formed using a sponge toxin (i.e a pore can be formed and then the pore collapses allowing the cell membrane integrity to be re-established). The term "pore" has been used to describe the

openings in the cell membranes produced by the sponge toxin. This seems appropriate when considering the transient changes in conductance observed, and channel-like events in artificial lipid bilayers. The advantage of reversibly producing a pore  
5 is that the pore can be used to introduce material, (e.g DNA or drugs) to a cell without permanently damaging the cell membrane, which can be lethal to the cell. This transient and stable transfection has not been shown by the prior art. The advantage of sponge toxins are that they are simple to use,  
10 highly chemically stable and have good water solubility. In this connection, in contrast to some methods of the prior art there is no need to encapsulate DNA in liposomes before transfection.

15 Preferably, the sponge toxin comprises at least one polymeric 1,3-alkylpyridinium salt (poly-APS). The halitoxin preparation mentioned herein is composed of many (tens) of distinct polymeric 1,3-alkylpyridinium compounds. This preparation has a mean molecular weight of about 5kDa. In  
20 contrast, the poly-APS preparation is composed of two polymeric 1,3-alkylpyridinium compounds with 29 and 99 repeating monomeric units as set out in Figure 1a.

Poly-APS has been shown in aqueous solutions to form non-  
25 covalently bound aggregates with a mean hydrodynamic radius of  $23 \pm 2$  nm. The radius of pores formed within these aggregates has been estimated on bovine erythrocytes by the use of osmotic protectants, and calculated with the Renkin equation to be about 2.9 nm. The combination of the pore sizes  
30 and the transient nature of the pore formation means that poly-APS is a preferable means of forming the reversible membrane pores, especially when macromolecules such as DNA are to be transported across the cell membrane.

Preferably, the sponge toxin is obtained from the sponge *Reniera sarai*, *Callyspongia ridleyi*, *Haliclona erina*, *Haliclona rubens*, *Haliclona viridis*, *Amphimedon viridis*,  
5 *Callyspongia fibrosa* and *Amphimedon compressa*.

Large numbers of sponge species produce pore-forming molecules. Halitoxin compounds have been isolated from a number of Haplosclerid genera.

10

Preferably, the sponge toxin has a molecular weight of between substantially 5 kDa and 20 kDa. Further preferably, the sponge toxin has a molecular weight of 5.5 kDa or 18.9 kDa. In a preferred embodiment, the composition comprises more  
15 than one sponge toxin each with a different molecular weight.

Conveniently the sponge toxin is formed from between 1 and 150 monomeric units as set out in Figure 1a. Preferably the sponge toxin is formed from between 20 and 100 monomeric  
20 units. In a preferred embodiment the sponge toxin is formed from 29 or 99 monomeric units.

Preferably, the concentration of sponge toxin is between 0.5 ng/ml and 5 µg/ml. Further preferably, the concentration  
25 of sponge toxin is between 0.5 ng/ml and 1 µg/ml. In a preferred embodiment the concentration of sponge toxin is between 0.5ng/ml and 0.5µg/ml.

According to a further aspect of the present invention,  
30 there is provided a method for the reversible formation of a membrane pore, the method comprising the steps of:-

a) incubating the membrane in the presence of a composition comprising sponge toxins; and

b) removing the composition from contact with the membrane.

This method of the present invention can be used in *in vitro* and *in vivo* techniques. In *in vitro* techniques, the present invention can be used as a biochemical research tool such as to study the transfection of cells with macromolecules such as proteins, DNA including plasmid cDNA, peptides, lipids, oligonucleotides and membrane impermeable such as fluorescent molecules (e.g. Lucifer yellow, neurobiotin, biocytins and dextran conjugates) and *in vivo* the method of the present invention can be used to transfer DNA to cells for gene therapy, in addition to the transfer of proteins, peptides, lipids and oligonucleotides.

15

Conveniently, a zinc, mercury, nickel or cobalt solution is used to attenuate the reversible formation of a membrane pore. Preferably, the concentration of the zinc solution is between substantially 1 to 2 mM. In preferred embodiments the concentration of zinc is 1.5 mM.

DNA, serum albumin and cholesterol can also be used to attenuate or prevent pore formation.

25 According to a yet further aspect of the present invention, there is provided a method for transfection of DNA into a cell *in vitro*, the method comprising the steps of:-  
a) incubating the cell in the presence of a composition comprising a sponge toxin;  
30 b) removing the composition from contact with the membrane;  
and  
c) adding nucleic acid.



The transfection of nucleic acid (e.g. DNA such as cDNA, RNA such as mRNA) into a cell in a stable manner is important as a biochemical research tool.

5        Preferably, the cell is incubated in the presence of composition for between 1 and 20 minutes prior to addition of nucleic acid. Further preferably, the cell is incubated in the presence of the composition for 5 minutes prior to the addition of nucleic acid.

10

It has been found that incubation for these periods results in an acceptable transfection rate with minimum disruption of the cell membrane.

15        Preferably, between 1.0 and 5.0 µg nucleic acid is added. Further preferably, 2.5 µg nucleic acid is added.

It has been found that the addition of these amounts of DNA result in an acceptable transfection rate. In this  
20 connection, an acceptable transfection rate is considered to be in the order of 20%.

Further preferably, the method comprises the additional steps of:- incubating the cell, in the presence of the  
25 composition and nucleic acid; and replacing the composition and DNA with a standard cell media.

Preferably, the cell is incubated in the presence of the composition and nucleic acid for between substantially 20 and  
30 200 minutes. Preferably, the incubation time is 180 minutes.

According to a further aspect of the present invention there is provided a method for transfection of a macromolecule

into a cell *in vivo*, the method comprising the step of:-

a) incubating the cell in the presence of a composition comprising a sponge toxin and the macromolecule.

5        Preferably, the macromolecule is cDNA, protein, peptide, lipid or oligonucleotide.

Further preferably, the macromolecule is the cytoskeletal protein tau.

10

Preferably, the cell is a hippocampal neurone.

According to a yet further aspect of the present invention, there is provided a model for use in the study of  
15 neurological disease or treatments thereof, the model comprising a rodent having undergone application of a composition comprising a sponge toxin, tau protein and phosphatase inhibitor to the hippocampus.

20        Preferably, the neurological disease is Alzheimer's.

Preferably, the rodent is a rat or a mouse.

According to further aspect of the present invention  
25 there is provided a method of studying a neurological disease, the method comprising:

a) applying a composition comprising a sponge toxin, tau protein and phosphatase inhibitor to the hippocampus of a rodent; and

30        b) studying the effect on the rodent.

Preferably, the phosphatase inhibitor is okadaic acid.

The invention will now be described in further detail with reference to the accompanying figures and experimental data, in which:-

5 Figure 1A shows the chemical structure of poly-APS from the marine sponge *Reniera sarai*. The preparation used was a mixture of polymers composed of repeating units (29 and 99 units).

10 Figure 1B shows the irreversible actions of poly-APS (50 µg/ml) on membrane potential and electrotonic potentials evoked by 100 pA hyperpolarising current commands.

Figure 1C shows the reversible actions of poly-APS (5 µg/ml)  
15 on membrane potential and electrotonic potentials (used to calculate input resistance). Records were obtained from cultured DRG neurones.

Figure 2A shows a bar chart showing the reversible actions of  
20 5 µg/ml poly-APS on membrane potential (n= 8). Poly-APS was applied to cultured DRG neurones for 20 seconds.

Figure 2B shows a bar chart showing the reversible actions of  
5 µg/ml poly-APS on input resistance (n= 8). Poly-APS was  
25 applied to cultured DRG neurones for 20 seconds.

Figure 3A shows a record of an irreversible rise in intracellular  $\text{Ca}^{2+}$ -evoked by 5 µg/ml poly-APS. These recordings are from a heterogeneous population of cells and  
30 so responses vary.

Figure 3B shows a record from a neurone in the same preparation as in Figure 3A showing a reversible rise in

intracellular  $\text{Ca}^{2+}$ -evoked by 5  $\mu\text{g/ml}$  poly-APS.

Figure 3C shows a trace from a single neurone showing the reversible action of 0.05  $\mu\text{g/ml}$  poly-APS.

5

Figure 4A shows the structure of halitoxin.

Figure 4B shows a record showing the irreversible collapse of membrane potential and input resistance evoked by 20 s application of 5  $\mu\text{g/ml}$  halitoxin. Top trace 50 pA hyperpolarising current commands evoked every 3 s, bottom trace showing the membrane potential and electrotonic potentials.

15 Figure 4C shows a record showing the reversible reductions in membrane potential and input resistance evoked by 20 s application of 5  $\mu\text{g/ml}$  poly-APS. Top trace shows 40 pA hyperpolarising current commands evoked every 3 s, bottom trace shows the membrane potential and electrotonic potentials.

25 Figure 4D shows a bar graph showing the effects of poly-APS (5  $\mu\text{g/ml}$ ) on membrane potential (filled bars) and input resistance (open bars), mean values are presented under control conditions (con;  $n = 9$ ), in the presence of toxin ( $n = 9$ ) and at 10 minutes recovery (rec;  $n = 6$ ).

Figure 5A shows a trace showing a single sustained response to the action of halitoxin preparations on intracellular  $\text{Ca}^{2+}$  in HEK293 cells loaded with fura-2. No responses were obtained with 0.005 and 0.05  $\mu\text{g/ml}$  halitoxin and \*\*\* denote the points of dye loss and cell damage produced by 5  $\mu\text{g/ml}$  halitoxin.

30

Figure 5B shows a trace showing a transient response to 0.5 µg/ml halitoxin. No responses were obtained with 0.005 and 0.05 µg/ml halitoxin and \*\*\* denote the points of dye loss and cell damage produced by 5 µg/ml halitoxin.

5

Figure 5C shows a single example trace showing the dose-dependent actions of poly-APS on fura-2 fluorescence ratio indicative of a rise in intracellular  $\text{Ca}^{2+}$ .

10 Figure 5D shows a single example trace showing transient and sustained responses to poly-APS (5 µg/ml) and halitoxin (5 µg/ml) respectively.

Figure 5E shows a bar chart showing the mean peak fluorescence  
15 values for experiments carried out to investigate the actions of 0.5 µg/ml poly-APS and halitoxin (open bars) and 5 µg/ml poly-APS and halitoxin (filled bars).

Figure 6A shows a record showing the lack of effect of acute  
20 application of NaCl-based extracellular solution containing 1.5 mM zinc applied after poly-APS had evoked changes in membrane potential and input resistance.

Figure 6B shows a record showing a slowed and reduced response  
25 to poly-APS after poly-APS was applied to the cells in the continual presence of NaCl-based extracellular solution containing 1.5 mM zinc. Constant current injection was applied at the end of the recording to hyperpolarise the membrane potential back to the resting level.

30

Figure 6C shows a bar chart showing the significant influence of the continued presence of zinc on poly-APS-evoked changes in membrane potential (n=9 and 7) and input resistance (n=8

and 7).

Figure 7A shows current voltage relationships for the poly-APS evoked currents in the presence and absence of zinc. Linear I/V relationships were obtained between 160 and 70 mV after leakage subtraction (the  $r^2$  values were 0.996 and 0.995 for poly-APS applied alone and in the continual presence of zinc respectively). Inset traces show currents evoked at 90 mV by poly-APS applied in the absence and presence of zinc.

10

Figure 7B shows a bar chart showing the inhibition of poly-APS (0.5  $\mu\text{g/ml}$ )-evoked inward currents by zinc. Control data show the mean holding currents required to clamp the neurones at 90 mV, zinc had no significant effect on the mean holding current. Poly-APS in the absence of zinc evoked a significant inward current ( $n=5$ ), however no significant inward current was produced by poly-APS in the presence of zinc ( $n=6$ ).

Figure 8A shows a record showing an increase in intracellular  $\text{Ca}^{2+}$  evoked by 5  $\mu\text{g/ml}$  poly-APS. Note that zinc was applied but that it was difficult to determine its actions due to the decay of the fluorescence ratio during washing. Irreversible responses were not attenuated by acute application of zinc.

25 Figure 8B shows a record showing the sustained but relatively modest rise in fluorescence ratio observed when poly-APS were applied in the continual presence of zinc.

Figure 8C shows a bar chart showing the mean imaging data derived from applying poly-APS (5  $\mu\text{g/ml}$ ) in the presence of zinc (filled bars,  $n=13$ ) and absence of zinc (open bars,  $n=46$ ). Zinc alone produced a significant but very modest increase in fluorescence ratio but subsequently attenuated the

response to poly-APS.

Figure 9A shows a record of fluorescence ratio changes produced in a single DRG neurone. Poly-APS (0.05 µg/ml) evoked a transient increase in  $\text{Ca}^{2+}$ ; zinc was then applied and caused a sustained rise in fluorescence. However, a further application of poly-APS (0.05 µg/ml) evoked no further increase in intracellular  $\text{Ca}^{2+}$  as reflected by the stable fluorescence ratio value.

10

Figure 9B shows a record of a similar experiment to that in Figure 9A except that zinc was not pre-applied but only simultaneously applied with the second application of poly-APS (0.05 µg/ml).

15

Figure 10A shows an example record from a HEK 293 cell showing that these cells did not respond to extracellular solution containing a depolarising concentration of KCl (30 mM) and therefore did not express voltage-activated  $\text{Ca}^{2+}$  channels.

20 However, this cell responded to the first application of poly-APS but showed only a modest increase in fluorescence ratio when poly-APS (0.5 µg/ml) was applied for a second time but with zinc present.

25 Figure 10B shows a bar chart showing mean fura-2 fluorescence ratios under control conditions, during stimulation with 30 mM KCl and during application of poly-APS without and with zinc (n=11).

30 Figure 11A shows HEK293 cells treated with lipofectamine, 0.5 µg/ml poly-APS or 0.5 µg/ml halitoxin in the absence (upper panels) or presence (lower panels) of pEGFP. Confocal images of fluorescent cells captured 48 h post-transfection, images

are representative of at least 3 other independent experimental repeats.

Figure 11B shows HEK293 cells transfected with pEGFP using 5 either lipofectamine or 0.05 µg/ml, 0.5 µg/ml, 5.0 µg/ml poly-APS/halitoxin were assessed for percentage of total cells fluorescent 48 h post-transfection. Data shown are means ± SEM, n = 3.

10 Figure 11C shows HEK293 cells transfected and assessed as in Figure 11B, which were subjected to crystal violet staining 48 h post-transfection to quantify intact and adherent cells. Data expressed as % of surviving cells relative to cDNA-only control, shown as means ± SEM, n = 3.

15

Figure 12A shows HEK293 cells co-transfected with pEGFP and pBABE cDNA using no transfection vehicle (left-hand panel), 0.5 µg/ml poly-APS (centre panel) or lipofectamine (right hand-panel). Following hygromycin B selection, cells were 20 imaged under confocal microscope and photographed. Photographs are representative of 3 other independent experimental repeats.

Figure 12B shows HEK293 cells co-transfected with TNFR2 and 25 pBABE cDNA using 0.5 µg/ml poly-APS were selected with hygromycin B and individual stable clones probed with no antibody (left-hand panels), TNFR1- (centre panels) or TNFR2-specific mouse monoclonal antibodies (right-hand panels). Subsequent secondary labelling with FITC anti-mouse IgG 30 allowed confocal visualisation. Shown here are untransfected HEK293 cells (upper panels), a positive TNFR2 clone (clone 4) (middle panels) and a negative TNFR2 clone (clone 7) (lower panels). Photographs are representative of 3 other independent



experiments.

Figure 12C shows the results from Poly-APS transfected stable TNFR2 clones which were dissociated, probed with no antibody, 5 TNFR1- or TNFR2-specific mouse monoclonal antibodies and secondary labelled with FITC anti-mouse IgG. Subsequent FACS analysis allowed quantitation and comparison of TNFR expression between untransfected HEK293 cells (left-hand panel), a TNFR2 positive clone (clone 4) (centre panel) and 10 a TNFR2 negative clone (clone 7) (right-hand panel). Histograms are representative of several other repeat experiments.

Figure 13 depicts two trails and the swimming paths taken by 15 the animals to the platform in the water maze experiments. Water maze experiments involve training animals to find a platform in a quadrant of a circular bath. Once the animal has learnt the position of the platform, the platform can be moved to test the animal's ability to show reversal learning. 20 With trial 1 the animal swims round the bath and has not learnt where the platform is, however during trial 16 the animal swims directly to the platform.

Figure 14A shows a bar chart showing that all animals learn 25 where the platform is in the water maze experiments, as reflected by the time (above 25%) spent in the quadrant with the platform.

Figure 14B shows in which quadrant of the circular bath the 30 platform was positioned during the water maze experiments for the above mentioned experiment identified in Figure 14A.

Figure 14C shows a line graph showing that all animals learn

where the platform is in the water maze experiments, as reflected by the decreasing swimming path lengths over a period of 4 days.

5 Figure 15A shows a bar chart showing that when the platform was moved into a different quadrant of the circular bath during the water maze experiments, the animals that have been treated with poly-APS, tau and okadaic acid fail to show learning reversal, while all other treatment groups do.

10

Figure 15B shows in which quadrant of the circular bath the platform was positioned during the water maze experiments for the above mentioned experiment identified in Figure 15A.

15 Figure 16 shows that brains from animals treated with poly-APS, tau and okadaic acid exhibit less long term potentiation (LTP). LTP is a form of neuronal plasticity which lasts many hours or days. It is also shown that if pathways get tetanic stimulation (arrow) subsequent electrophysiological signalling  
20 is increased. The experiments were performed *ex vivo* on hippocampal slices.

Figure 17 shows the delivery of tau to the intracellular compartments of hippocampal neurones in culture. The tau  
25 levels (black) are detected using tau-specific antibodies.

Figure 18 shows a record of action potentials from cultured hippocampal neurones treated with poly-APS and tau. Neurones treated with poly-APS, tau and okadaic acid fail to show  
30 action potential firing even when stimulated with much larger currents.

Figure 19 shows a record from cultured hippocampal neurones

stimulated with KCl pulses showing that neurones treated with poly-APS, tau and okadaic acid fail to show proper calcium transients. Calcium transients were measured with fura-2 calcium imaging. Combined with the data recorded in Figure 5 18, this shows that the excitability of these neurones is inhibited, probably by mis-functioning of voltage-activated channels.

Figure 20 shows a record of mean data for calcium transients 10 exhibited by neurones exposed to the different treatments identified thereon. Neurones treated with poly-APS, tau and okadaic acid clearly exhibit a distinct fluorescence ratio. Combined with the data recorded in Figures 18 and 19, this shows that the excitability of these neurones is greatly 15 reduced, probably involving mis-functioning of voltage-activated channels.

#### Example 1

#### 20 Sponge toxin preparations

Poly-APS (Fig. 1A) were purified from the marine sponge Reniera (=Haliclona) sarai Pulitzer-Finali (Halicionidae). Aqueous extract containing poly-APS was passed through an 25 ultrafiltration membrane (3 kDa cut off). Toxins were eluted from Sephadex G-50 and Sephacryl S-200 (Sepčić et al 1997; J. Nat. Prod. 60 991-996). A halitoxin preparation was isolated from *Callyspongia ridleyi*. Butanol extraction containing halitoxins was applied to a lipophilic Sephadex LH-20 size 30 exclusion column and the 5 to 6 kDa fraction isolated and studied (Jaspers et al 1994; J. Org. Chem. 59 3253-3255 and Scott et al 2000; J. Mem. Biol. 176 119-131). Stock solutions containing 5 mg/ml of either toxin preparation were diluted

to produce poly-APS or halitoxin test solutions.

#### Cell culture

5        Primary cultures of dorsal root ganglion (DRG) neurones  
were prepared following enzymatic and mechanical dissociation  
of dorsal root ganglia from decapitated 2-day old Sprague  
Dawley rats. The sensory neurones were plated on laminin-  
polyornithine coated coverslips and bathed in F14 culture  
10 medium (Imperial Laboratories) supplemented with 10% horse  
serum (Gibco), penicillin (5000 IU/ml), streptomycin (5000  
mg/ml),  $\text{NaHCO}_3$  (14 mM) and nerve growth factor (20 ng/ml). The  
cultures were maintained for up to 2 weeks at 37°C in  
humidified air with 5%  $\text{CO}_2$ , and re-fed with fresh culture  
15 medium every 5-7 days.

HEK293 cells (human embryonic kidney cell line) were  
maintained in Dulbecco's modified Eagle's medium (DMEM). In  
preparation for transfection experiments cells were seeded 24  
20 hours prior to experimentation at a density of  $5 \times 10^5$  cells  
per well in six well plates, the same seeding density was used  
for confocal and electrophysiology/calcium imaging experiments  
in 35 mm culture dishes.

#### 25 Electrophysiology and calcium imaging.

All experiments were conducted at room temperature  
(approximately 23°C). The whole-cell recording technique was  
used to study the actions of poly-APS and halitoxin  
30 preparations on membrane potential, input resistance  
(evaluated from 100-300 ms electrotonic potentials evoked by  
-30 to -200 pA current commands) and holding current. Whole  
cell recordings were made using an Axoclamp-2A switching

voltage clamp amplifier operated at a sampling rate of 15-20 kHz. Low resistance (4-10 or 4-7 M $\Omega$ ) borosilicate glass patch pipettes were fabricated using a Kopf model 730, needle/pipette puller. The neurones were bathed in a NaCl-based extracellular solution containing in mM: NaCl, 130; KCl, 3.0; CaCl<sub>2</sub>, 2.0; MgCl<sub>2</sub>, 0.6; NaHCO<sub>3</sub>, 1.0, HEPES 10.0, glucose 5.0. NaCl-based extracellular solution with zinc was also made up separately and contained 1.5 mM zinc (zinc atomic absorption standard solution, Sigma). The pH and osmolarity of extracellular solutions were adjusted to 7.4 and 310-320 mOsmol/l with NaOH and sucrose respectively. The patch pipette solution contained in mM: KCl, 140; EGTA, 5; CaCl<sub>2</sub>, 0.1; MgCl<sub>2</sub>, 2.0; HEPES, 10.0; ATP, 2.0; and the pH and osmolarity were adjusted to 7.2 with Tris and 310-315 mOsmol/l with sucrose. For voltage clamp recordings, neurones were held at -90 mV and linear current-voltage relationships were generated with 100 ms voltage step commands to potentials between -170 and -60 mV. Poly-APS and halitoxin were applied to the extracellular environment by low pressure ejection via a blunt micropipette (tip diameter about 10  $\mu$ m) positioned approximately 100  $\mu$ m from the neurone being recorded. The cells were maintained in a bath and were not continually perfused, drug concentrations declined after pressure ejection as a result of diffusion.

25

The electrophysiological data were stored on digital audio tape (DAT) using a DTR-1200 DAT recorder (Biologic) and subsequently analysed using Cambridge Electronic Design voltage clamp software (version 6). For monitoring changes in membrane potential or holding current continuous records were obtained on a chart recorder (Gould 2200s pen recorder).

Cultured DRG neurones and HEK 293 cells were incubated

for 1 hour in NaCl-based extracellular solution containing 10  $\mu$ M fura-2AM (Sigma, 1mM stock in dimethylformamide). The cells were then washed for 20 minutes to remove the extracellular fura-2AM and to allow cytoplasmic de-esterification of the 5  $\text{Ca}^{2+}$  sensitive fluorescent dye. The cells were constantly perfused (1 - 2 ml/min) and viewed under an inverted Olympus BX50W1 microscope with a KAI-1001 S/N 5B7890-4201 Olympus camera attached. The fluorescence ratiometric images from the data obtained at excitation wavelengths of 340nm and 380nm 10 were viewed and analysed using OraCal pro, Merlin morphometry temporal mode (Life Sciences resources, version 1.20).

All data are given as mean  $\pm$  standard error of the mean (SEM) and statistical significance was determined, using the 15 Student's two-tailed t test, paired or independent where appropriate and P values are reported in the text.

#### DNA transfer

20 Plasmid cDNAs used were pEGFP-C1 (Clontech), an enhanced green fluorescent protein (EGFP) cDNA vector under control of a constitutively active SV40 promoter (pEGFP), human tumour necrosis factor receptor-2 (TNFR2) cDNA (provided by Werner Lesslauer, Basel, Switzerland) and pBABE hygromycin resistance 25 cDNA (Stratagene). Control transfections were carried out using optimized lipofectamine (Invitrogen life technologies) lipid-micelle-mediated transfection protocol, which incubates cells with 1  $\mu$ g cDNA and lipofectamine in the absence of serum for 3 h prior to re-introduction to serum-containing medium. 30 A standard toxin transfection protocol was developed and optimised throughout the passage of this work. The protocol involved a 5 min serum-free cell incubation with 0.5  $\mu$ g/ml of a sponge toxin preparation, followed by addition of 2.5  $\mu$ g

cdNA. After a further 3 hour incubation, DMEM medium was replaced with standard serum-containing medium. Lipofectamine and poly-APS stable transfections involved co-transfection with 1.0 µg or 2.5 µg cdNA respectively (either pEGFP or TNFR2 5 cdNA) and 1.0 µg of pBABE cdNA. Colonies of stably transfected cells were selected in DMEM containing 100 µg/ml hygromycin B (Boehringer Mannheim). Once established, colonies were harvested using cloning discs and trypsin-EDTA and seeded into larger vessels until sufficient cells were available for 10 analysis. In order to reduce variability cdNA was introduced into 1 ml preparations at a volume of 10 µl.

#### Crystal violet assay

15 Adherent cells were fixed in paraformaldehyde and stained in crystal violet dye (Sigma), subsequent elution and spectrophotometric analysis quantified the amount of intact viable cells following transfection.

#### 20 EGFP detection

Cellular expression of EGFP was evaluated using an Eppendorf fluorescence/visible light microscope set-up to directly assess the percentage of total cells fluorescing. 25 Additionally, cells were imaged using a BioRad pradiance confocal system.

#### FACS analysis

30 HEK293 cells stably over-expressing TNFR2 were dissociated from their culture vessels with 3 ml cell dissociation solution (trypsin free, Sigma) and assessed for TNFR1 and TNFR2 expression with specific mouse mAb (htr-9 and

utr-1 respectively), secondary labelling with flouoroisothiocyanate (FITC) anti-mouse IgG followed by Fluorescence-activated cell sorting analysis (Becton-Dickenson)

5

#### Confocal analysis

HEK293-TNFR2 stable clones were plated into triplicate  
10 35 mm dishes and fixed with ice-cold methanol. Individual dishes were probed with no antibody, TNFR1- or TNFR2-specific mouse mAb (htr-9 and utr-1 respectively) and then secondary labeled with FITC anti-mouse IgG. Cellular FITC labeling was then assessed using a BioRad pradiance confocal system, and  
15 was indicative of TNF receptor expression levels.

Delivery of the cytoskeletal protein tau and poly-APS into the rat hippocampus

20 This was performed using well known methods of the art. The tau protein plus poly-APS, as appropriate, were injected into the hippocampus through a bore hole. The concentration of poly-APS used was between 0.005 µg/ml and 5 µg/ml, preferably 0.005 µg/ml or 5 µg/ml. The concentration of tau  
25 was 1 µg/ml. Okadaic acid was administered via a mini pump at a concentration of 1.2nM for 7 days.

Obtaining ex vivo hippocampal slices.

30

#### Recordings

Hippocampal slices were obtained from pigmented Lister-



Hooded rats 3-4 weeks after birth. Methods have been described in detail previously (White and Platt (2000) Eur J Neurosci 12 3155-3162). Briefly, terminally anaesthetised animals were decapitated, the brain quickly removed and placed into ice-cold artificial cerebrospinal fluid (aCSF, See Table 3 for composition) saturated with 95% O<sub>2</sub>/5% CO<sub>2</sub>, pH 7.4. A vibrotome (Oxford Instruments, UK) was used to prepare 350- $\mu$ m parasagittal hippocampal slices, which were subsequently maintained in aCSF at ca 30°C for at least an hour before recordings commenced.

#### Measurement of Long-Term Potentiation

For extracellular field recordings, slices were submerged in a recording chamber; a monopolar stimulation electrode (ca 0.5 M $\Omega$ , WPI, USA) driven by a DS2 A stimulus isolator (Digitimer, UK) was placed in the outermost part of the optic layer and a recording glass electrode (Clark, UK) filled with aCSF was placed in the superficial grey layer adjacent the stimulation electrode. Evoked glutamate-mediated fEPSPs were recorded (Axoclamp 2A, bridge mode; Axon, USA), stored on a computer and analysed (amplitude and slope of the initial rising phase, from fibre volley to peak; PWIN software package). Activity was also continuously monitored on an oscilloscope (Tektronix, USA). As in previous papers (Platt and Withington (1998) Neuropharmacology 37 1111 - 1122; White and Platt (2000) Eur J Neurosci 12 3155-3162), slices were stimulated every 30 s at 70-80% of the maximum fEPSPs until a stable baseline was obtained for at least 20 min.

30

#### Drugs

LTP<sub>c</sub> was induced by superfusion of 3 mM GABA for 7 min

and was washed out with aCSF for another 60 min. In subsequent experiments the perfusion solution was manipulated by either decreasing potassium ions, calcium ions, or by replacing bicarbonate buffer with *N*-[hydroxyethyl] piperazine-5 *N*-[2-ethanesulfonic acid] (HEPES; see Table 3 for details of aCSF composition). The pH was monitored repetitively, in particular for experiments in HEPES-buffered solution, and found to be stable throughout the experiments. A CA inhibitor ethoxyszolamide (EZA; 50  $\mu$ M) was also used. NKCC1 and KCC2 inhibitors bumetanide (50  $\mu$ M) and furosemide (100  $\mu$ M) were also tested. All drugs were purchased from Sigma Aldrich, USA apart from the salts used for aCSF which were obtained from BDH, UK.

Table 3 Composition of extracellular solutions (artificial cerebrospinal fluid) in Mm

Compound	Control	Low K <sup>+</sup>	Low Ca <sup>2+</sup>	HEPES
NaCl	127	129.5	130	127
20 KCl	4	1.5	4	4
CaCl <sub>2</sub>	2.5	2.5	1	2.5
KH <sub>2</sub> PO <sub>4</sub>	1.5	1.5	1.5	1.5
MgCl <sub>2</sub>	-	-	-	-
MgSO <sub>4</sub> .7H <sub>2</sub> O	1.3	1.3	3	1.3
25 Glucose	10	10	10	10
NaHCO <sub>3</sub>	25	25	25	-
HEPES	-	-	-	10

#### Data Analysis

30

LTP<sub>6</sub> was calculated relative to baseline values in percent for both slope and amplitude. Statistical significance

was assessed for data before and 1 h after GABA application (three consecutive values were averaged in each case) using Prism for Windows (Version 3.00; GraphPad Software, USA). One-way ANOVA with Dunnett's multiple comparison *post hoc* test was performed for LTP<sub>G</sub> comparison 60 min after GABA application. Unpaired and paired two-tailed Student's *t*-tests were also used to compare LTP<sub>G</sub> in different conditions and to compare baseline and LTP<sub>G</sub> values within groups, respectively.

## 10 The Open-field Water Maze For Rats

### Apparatus

The open-field water maze used has been described previously. In brief, a pool 2 m in diameter was filled with water at a temperature of  $25 \pm 2^{\circ}\text{C}$ , to a depth of approximately 50 cm, and rendered opaque by the addition of latex subflooring compound. A platform, 11 cm in diameter, submerged approximately 2 cm below the water surface, provided the only means of escape for the rats. The pool was located in a room that provided plenty of extramaze cues. A curtain could be drawn around the pool's circumference.

### Behavioural Paradigm

During habituation and all subsequent training and testing, the following conditions applied. Four cardinal points (north, south, east and west) were semirandomly chosen as start locations, and rats were released facing the wall of the pool. If the rats could not locate a platform within 120 s, the trial was terminated and the rat was guided to the platform by the experimenter. After having found the platform, rats were left on it for 30 s before being placed back in their cage. Heating lamps placed in a room adjacent the water maze room were used to dry the rats after testing.

## Results

### Irreversible and reversible actions of poly-APS on 5 electrophysiological properties of cultured DRG neurones

Extracellular application of 50  $\mu\text{g/ml}$  poly-APS for approximately 20 seconds resulted in an irreversible collapse in membrane potential and a dramatic fall in input resistance  
10 in all DRG neurones studied (Fig. 1B). The mean membrane potential was depolarised from  $-65 \pm 1 \text{ mV}$  to  $-9 \pm 3 \text{ mV}$  ( $n=17$ ;  $P<0.0001$ ) and this was associated with a reduction in input resistance from  $350 \pm 79 \Omega$  to  $56 \pm 27 \Omega$  ( $n=17$ ;  $P<0.0001$ ). No recovery was seen 120 minutes after removal of the perfusion  
15 pipette containing Poly-APS. At concentrations of 5 and 0.5  $\mu\text{g/ml}$ , poly-APS evoked reversible depolarisations and associated decreases in input resistance (Fig. 1C). Poly-APS showed a degree of dose dependent action when the sponge toxins were applied for approximately 20 seconds, the mean  
20 percentage reductions in input resistance were 84%, 67% and 17% for 50, 5, and 0.5  $\mu\text{g/ml}$  poly-APS respectively. Recovery periods varied but were seen within 20 minutes of removing the perfusion pipette containing poly-APS. In responses to 5  $\mu\text{g/ml}$  poly-APS, 6 out of 8 neurones showed at least partial recovery  
25 of membrane potential and input resistance after toxin application (Figure 2A and figure 2B).

### Reversible actions of poly-APS on cell membrane calcium permeability

30

Consistent with the electrophysiological results,  $\text{Ca}^{2+}$  imaging experiments on cultured DRG neurones showed that poly-APS evoked irreversible (Figure 3B) and reversible (Figure 3B)

increases in intracellular  $\text{Ca}^{2+}$ . Although the responses were very variable a dose-dependent trend was clearly apparent. Poly-APS (5  $\mu\text{g/ml}$ ) produced mainly responses that either only partially recovered or were irreversible and 0.05  $\mu\text{g/ml}$  poly-APS evoked predominantly transient responses with 4 out of 45 neurones failing to respond (Table 1). These results highlight the narrow concentration range required for reversible pore formation and the reason for cell death at high concentrations of poly-APS.

10

Table 1. Reversible and irreversible rises in intracellular  $\text{Ca}^{2+}$ -evoked by poly-APS as detected using the  $\text{Ca}^{2+}$ -sensitive ratiometric dye fura-2. Reversible is defined as a transient event that showed recovery of 75% or more. Irreversible is defined as a sustained event that only decayed by 10% or less during washing.

20	Poly-APS Concentration ( $\mu\text{g/ml}$ )	Irreversible (I) or Reversible (R) responses	Amplitude of response in ratio units $\pm$ S.E.M.	Number of DRG neurones	Percentage of DRG neurones
	5*	I	$2.54 \pm 0.28$	15	29%
	5*	R	$2.18 \pm 0.45$	9	17%
	0.5	I	3.16	1	3%
25	0.5	R	$2.58 \pm 0.51$	31	97%
	0.05	I	-	0	0%
	0.05	R	$2.53 \pm 0.4$	41	100%

\*28 neurones (54%) showed partial recovery and were not included in this analysis.

No significant differences were observed when comparing the mean amplitudes of irreversible and reversible rises in intracellular  $\text{Ca}^{2+}$  and the mean amplitude of events evoked by 5, 0.5 and 0.05  $\mu\text{g/ml}$  poly-APS. This relates partially to the

considerable variability of responses, for example the maximum and minimum changes in fluorescence ratio produced by 5 µg/ml poly-APS were 5.42 and 0.48, and even in neighbouring neurones during the same experiment great differences in responses were 5 seen. Additionally, poly-APS was applied for 2 to 2.5 minutes, which was significantly longer than the periods the toxins were applied for in the electrophysiological experiments. One benefit of using cultured DRG neurones in this study was that these neurones express voltage-activated  $\text{Ca}^{2+}$  channels, which 10 can be used to assess the viability of neurones after exposure to poly-APS. The experimental trace in Figure 3C shows that  $\text{Ca}^{2+}$  transients evoked by a depolarising pulse of extracellular solution containing KCl (30 mM) could be produced before and after application of 0.05 µg/ml poly-APS. 15 Accordingly, it was shown that repeatable poration could be obtained and entry of  $\text{Ca}^{2+}$  via voltage-gated channels remained intact after application of low doses of poly-APS indicating that at least at low concentrations cytotoxic damage did not occur as a result of pore formation. It was however surprising 20 that voltage-gated  $\text{Ca}^{2+}$  channels still functioned after a period of poration given the sensitivity of  $\text{Ca}^{2+}$  channels to intracellular  $\text{Ca}^{2+}$ -induced inactivation.

Action of poly-APS and Halitoxin preparations on membrane 25 potential, input resistance and  $\text{Ca}^{2+}$  permeability of HEK293 cells

The actions of poly-APS and the halitoxin preparations on membrane potential, input resistance and  $\text{Ca}^{2+}$  permeability 30 of HEK293 cells were also evaluated. A 20 second application of halitoxin (0.5 and 5.0 µg/ml) predominantly resulted in irreversible effects on the electrophysiological properties of HEK293 cells (Fig. 4B). Halitoxin (0.5 µg/ml) reduced

membrane potential from  $-45 \pm 5\text{mV}$  to  $-16 \pm 5\text{mV}$  ( $n = 6$ ;  $P < 0.003$ ) and partial but significant recovery to  $-26 \pm 4\text{mV}$  ( $n = 6$ ;  $P < 0.02$ ) was observed 10-20 min after toxin application. Halitoxin ( $5 \mu\text{g/ml}$ ) reduced membrane potential from  $-46 \pm 3\text{mV}$  to  $-3 \pm 1\text{mV}$  ( $n = 5$ ;  $P < 0.003$ ) with no significant recovery. Similar trends were observed when input resistance was determined from the electrotonic potentials evoked by hyperpolarising current commands. For example the resting input resistance was reduced from  $903 \pm 313 \text{ M}\Omega$  to  $315 \pm 206 \text{ M}\Omega$  ( $n = 6$ ;  $P < 0.04$ ) by  $0.5 \mu\text{g/ml}$  halitoxin but no significant partial recovery was observed. There was also evidence that the effects on membrane potential were apparently dose dependent with  $0.5$  and  $5 \mu\text{g/ml}$  halitoxin reducing membrane potentials by  $64 \pm 10\%$  ( $n = 6$ ) and  $94 \pm 3\%$  ( $n = 5$ ;  $P < 0.03$ ) respectively. In contrast the actions of poly-APS applied for 20 seconds were predominantly reversible both at  $5 \mu\text{g/ml}$  ( $n = 6$  out of 9 cells, Figure 4C and Figure 4D) and at  $0.5 \mu\text{g/ml}$  ( $n = 6$  out of 9 cells). The apparent dose dependency was less apparent and both  $5$  and  $0.5 \mu\text{g/ml}$  poly-APS gave equivalent electrophysiological responses to those obtained with  $0.5 \mu\text{g/ml}$  halitoxin.

Experiments using the ratiometric dye fura-2 also indicated that both halitoxin and poly-APS evoked large changes in intracellular  $\text{Ca}^{2+}$ . Initially a protocol was used to examine the dose-dependent effects of the sponge toxins. Halitoxin or poly-APS were applied for 140 seconds at doses of  $0.005$ ,  $0.05$ ,  $0.5$  and  $5.0 \mu\text{g/ml}$  with variable wash periods between each application. Few cells responded to the sponge toxins when they were applied for 140 seconds at doses of  $0.005 \mu\text{g/ml}$  (10 of 115 cells for Halitoxin and 2 of 90 cells for poly-APS). Accordingly,  $0.005 \mu\text{g/ml}$  appears to be the threshold concentration for a detectable  $\text{Ca}^{2+}$  transient in

HEK293 cells. The halitoxin preparation produced some anomalous results because cell lysis and dye loss were usually observed with 5  $\mu\text{g/ml}$  when the cells had been exposed previously to the three lower doses of toxin. The mean peak 5 changes in fluorescence ratios observed were  $0.2 \pm 0.03$  ( $n = 10$ ),  $0.43 \pm 0.03$  ( $n = 80$ ) and  $1.09 \pm 0.04$  ( $n = 114$ ) with 0.005, 0.05 and 0.5  $\mu\text{g/ml}$  halitoxin respectively. However, at least in some cells, transient changes in intracellular  $\text{Ca}^{2+}$  were observed with 0.05 and 0.5  $\mu\text{g/ml}$  halitoxin (Fig. 5A).

10 Dose-dependent and predominantly reversible responses were obtained with poly-APS (0.005 - 5  $\mu\text{g/ml}$ ; Fig. 5B). The mean peak changes in fluorescence ratios observed were 0.1 ( $n = 2$ ),  $0.65 \pm 0.06$  ( $n = 63$ ),  $0.88 \pm 0.06$  ( $n = 83$ ) and  $1.17 \pm 0.05$  ( $n = 89$ ) with 0.005, 0.05, 0.5 and 5.0  $\mu\text{g/ml}$  poly-APS

15 respectively. The different  $n$  numbers for both sets of data are indicative of varied threshold sensitivities of different HEK293 cells to the sponge toxins. In this connection, some cells did not respond to 0.05  $\mu\text{g/ml}$  toxin. Separately applying single doses of each toxin clearly showed the different

20 amplitudes and natures of responses produced by the sponge toxins (Figure 5C and Figure 5D). All cells studied responded to 0.5 and 5.0  $\mu\text{g/ml}$  toxins and, therefore, these doses were used in subsequent experiments.

## 25 Actions of zinc on pore formation by poly-APS.

The hemolytic actions of poly-APS have been found to be attenuated by zinc, most likely in the ionic form  $\text{Zn}^{2+}$ , so the protective properties of zinc ( $\text{Zn}^{2+}$ ) when applied continually

30 or after pore formation by poly-APS was investigated. Experiments were carried out on the same cultures of DRG neurones to allow comparisons between results obtained using the two protocols. Using the first experimental protocol,



poly-APS (50  $\mu\text{g/ml}$ ) reduced the membrane potential from  $-66 \pm 2 \text{ mV}$  to  $-13 \pm 6 \text{ mV}$  ( $P < 0.0001$ ) but during subsequent application of extracellular solution containing 1.5mM zinc the membrane potential did not significantly recover and had a value of  $-17 \pm 6 \text{ mV}$  ( $n=8$ ). A similar pattern of results was obtained for input resistance measurements which had values of  $252 \pm 52 \Omega$ ,  $13 \pm 3 \Omega$  (significance compared to control  $P < 0.005$ ) and  $19 \pm 7 \Omega$  ( $n=8$ ), under control conditions, during application of 50  $\mu\text{g/ml}$  poly-APS and during application of zinc. It was clear from this experiment that once poly-APS had produced pores or lesions in the cell membrane zinc failed to attenuate the conductances (Fig. 6A). However, using the second experimental protocol which involved continually bathing neurones with extracellular solution containing 1.5 mM zinc resulted in significant inhibition of poly-APS actions (Figure 6B and Figure 6C). In the presence of zinc the resting membrane potential was  $-47 \pm 7 \text{ mV}$  and application of poly-APS (50  $\mu\text{g/ml}$ ) with zinc significantly reduced the membrane potential to  $-27 \pm 6 \text{ mV}$  ( $n=7$ ;  $P < 0.07$ ). Similarly in the presence of zinc the input resistance was  $261 \pm 34 \Omega$  and application of poly-APS (50  $\mu\text{g/ml}$ ) with zinc significantly reduced the input resistance to  $114 \pm 62 \Omega$  ( $n=7$ ;  $P < 0.05$ ). Figure 6C shows a bar chart of data normalised with respect to the resting values to illustrate the significant attenuation of poly-APS effects when zinc was present prior to and during application of the toxin.

Experiments were then conducted under voltage clamp conditions to remove the influence of voltage-activated channels. Neurones were held at  $-90 \text{ mV}$  and currents were activated by poly-APS (10  $\mu\text{g/ml}$ ) in the presence and absence of 1.5 mM zinc containing solution. Current voltage relationships were generated between  $-160 \text{ mV}$  and  $-70 \text{ mV}$  and

subtracted to produce difference current voltage relationships for the poly-APS evoked currents (Figure 7B). The results with 10  $\mu\text{g/ml}$  poly-APS showed that although zinc significantly reduced the action of the toxin its effects were not abolished 5 so the concentration of poly-APS was reduced. The presence of 1.5 mM zinc containing solution prevented significant activation of inward currents evoked by 0.5  $\mu\text{g/ml}$  poly-APS. In the absence of zinc the holding current required to clamp the cell membrane at -90 mV increased from  $-150 \pm 30$  pA to 10  $-600 \pm 190$  pA ( $n=5$ ;  $P<0.04$ ) during application of 0.5  $\mu\text{g/ml}$  poly-APS. In contrast with 1.5 mM zinc containing solution continually present, the holding current was not significantly increased by poly-APS (0.5  $\mu\text{g/ml}$ ) from a resting level  $-120 \pm 30$  pA to  $-170 \pm 70$  pA ( $n=6$ ; NS). The mean amplitudes of 15 currents evoked by poly-APS in the presence and absence of zinc were also significantly different ( $P<0.05$ ; Fig. 7B).

Imaging experiments were then carried out to determine whether zinc would attenuate the rise in  $\text{Ca}^{2+}$  evoked by poly- 20 APS. Protocols were again designed to assess the ability of zinc to inhibit rises in  $\text{Ca}^{2+}$  produced by poly-APS, when applied before or after poration of the cell membrane. It was not possible to clearly characterise the effects of zinc when applied after poly-APS because of the slow partial recovery 25 of the responses that occurred during washing (Fig. 8A). Application of zinc itself produced a modest but significant increase in fluorescence ratios above basal levels, and poly-APS (5  $\mu\text{g/ml}$ ) applied with zinc subsequently evoked a sustained rise in fluorescence ratio (Fig. 8B). However, these 30 responses in the presence of zinc were significantly smaller than the responses evoked by poly-APS in the absence of zinc (Figures 8A, 8B & 8C). The change in the mean fluorescence ratios evoked by poly-APS in the absence and presence of zinc

were significantly different and had values of  $3.46 \pm 0.18$  (n=46) and  $1.2 \pm 0.05$  (n=13;  $P < 0.0001$ ) respectively. These experiments were conducted on two populations of neurones; however, it was hoped that by lowering the poly-APS concentration it would be possible to record poly-APS responses in the absence and presence of zinc in the same neurones. Preliminary observations showed that repeatable responses to poly-APS could be obtained from some neurones. In these experiments the mean control fluorescence ratio was  $0.9 \pm 0.03$ . The first and second applications of poly-APS ( $0.5 \mu\text{g/ml}$ ) resulted in significant increases in fluorescence ratio to values of  $5.36 \pm 1.38$  and  $3.45 \pm 0.55$  (n=10;  $P < 0.01$  and  $0.05$  compared to control). The first and second responses to poly-APS were not significantly different.

15

Subsequently experiments were conducted with  $0.05 \mu\text{g/ml}$  poly-APS. For this experiment the mean fluorescence ratio under control conditions was  $0.94 \pm 0.02$ . The first response to poly-APS ( $0.05 \mu\text{g/ml}$ ) resulted in a significant rise in fluorescence ratio to  $2.53 \pm 0.31$  (n=22;  $P < 0.01$ ); subsequent application of zinc and a second application of poly-APS ( $0.05 \mu\text{g/ml}$ ) produced an attenuated response as the fluorescence ratio increased to  $1.85 \pm 0.16$  (n=22). The second response in the presence of zinc was significantly different from both the basal levels ( $P < 0.01$ ) and the first response ( $P < 0.05$ ). The second response is likely to be an overestimation of the poly-APS actions because zinc itself produced more pronounced effects than predicted (Fig. 9A). Similar responses were obtained when poly-APS and zinc were simultaneously applied to produce the second response (Fig. 9B). It is likely that zinc passed through residual pores in the membrane produced during the first application of poly-APS and resulted in a slow sustained increase in fluorescence ratio.

Actions of zinc on pore formation by poly-APS in HEK 293 cells.

5        One difficulty of assessing rises in intracellular  $\text{Ca}^{2+}$  evoked by poly-APS in cultured DRG neurones is that in addition to  $\text{Ca}^{2+}$  entry through pores formed by poly-APS, rises in intracellular  $\text{Ca}^{2+}$  will also be produced by activation of voltage-gated  $\text{Ca}^{2+}$  channels and mobilisation of  $\text{Ca}^{2+}$  from  
10 stores. Although it should be noted that repeatable responses to low doses of poly-APS were obtained, it is possible to underestimate or overestimate the inhibitory effects of zinc given the other pathways for raising intracellular  $\text{Ca}^{2+}$ . For this reason preliminary experiments were conducted on HEK 293  
15 cells which did not express voltage-gated  $\text{Ca}^{2+}$  channels. As with the DRG neurones, poly-APS (1.5  $\mu\text{g}/\text{ml}$ ) evoked repeatable rises in intracellular  $\text{Ca}^{2+}$ . The mean fluorescence ratios for the first and second applications of poly-APS were  $2.31 \pm 0.25$  and  $1.96 \pm 0.3$  ( $n=7$ ; not significantly different).

20

Experiments were then carried out to assess the influence of zinc on the  $\text{Ca}^{2+}$  entry produced by poly-APS. Under control conditions the mean fluorescence ratio was  $1.16 \pm 0.07$  and brief application of extracellular recording solution  
25 containing 30 mM KCl caused no significant response ( $n=11$ ; Fig. 10A & 10B). Application of poly-APS (0.5  $\mu\text{g}/\text{ml}$ ) in the absence and presence of zinc evoked responses, and it was clear from these results that zinc attenuated the rise in intracellular  $\text{Ca}^{2+}$  produced by poly-APS.

30

Thus, it was clearly demonstrated that zinc ( $\text{Zn}^{2+}$ ), if continually present during application of poly-APS, could attenuate the effects of the sponge toxin preparation on

membrane potential, input resistance, whole cell currents and  $\text{Ca}^{2+}$  permeability.

Further, the results show that poly-APS can reversibly  
5 form pores in the membranes of DRG neurones and HEK 293 cells,  
and that applying extracellular zinc with the preparation of  
sponge toxins can attenuate these actions. The survival of  
cells even after dramatic but temporary changes in membrane  
potential and input conductances means that transient pore  
10 formation by low doses of poly-APS could be used to deliver  
materials to the intracellular environment, without cell  
damage.

Previous studies have been performed on pore properties  
15 and potential mechanisms of passive DNA flow. Furthermore  
single stranded polynucleotide RNA and DNA molecules have been  
shown to traverse lipid bilayers treated with *Staphylococcus*  
*aureus*  $\alpha$ -toxin or  $\alpha$ -hemolysin, which creates membrane pores  
of 1 - 2.5 nm diameter. The passage of these nucleotide  
20 molecules across the bilayer was signalled by fluctuations in  
channel conductances as individual molecules passed through,  
and allowed molecular characterisation. A number of other  
diverse molecules have been similarly shown to create  
perforations in lipid bilayers, including perforin,  
25 complement, detergents such as  $\alpha$ -escin and saponin, and  
alamethicin but as yet none has been shown to provide passage  
for DNA. More conventional approaches to DNA transfer across  
lipid bilayers and into intact cells include lipid-micelle-  
mediated transfection, e.g. lipofectamine, electroporation and  
30 micro-injection, but these approaches yield variable results  
and can be difficult to control, often requiring specialist  
equipment or laborious pretreatments and optimisation.

Poly-APS has distinct but comparable properties with cationic surfactants but variation in the extent of polymerisation of the poly-APS appears to modulate the biological activity in an unpredictable manner. In the present  
5 invention we have identified reversible effects of poly-APS on cell membrane properties. Previously, the related but lower molecular weight preparation of halitoxins evoked irreversible pore formation and associated changes in cell membrane input conductance. There are several possible explanations for the  
10 reversible changes in membrane potential, input resistance, currents and intracellular  $\text{Ca}^{2+}$  seen with poly-APS. Firstly, the larger degree of polymerisation in poly-APS, relative to smaller molecules composed of fewer monomeric units, may result in less stable interactions in the cell membranes and  
15 allow wash out of the larger pore forming sponge toxins with a resulting temporary pore formation. Secondly, and perhaps more likely is the possibility that the larger poly-APS may be sufficiently flexible so that lipids can re-arrange themselves after pore formation and thus block the ion  
20 conducting pathways through the cell membrane. This may occur due to "hydrophobic collapse" when the toxin is in the membrane. In such a circumstance, alkyl chains may surround all the pyridinium groups so that they are compatible with the membrane. The ability of poly-APS to form non-covalently bound  
25 aggregates with a mean hydrodynamic radius of 23 nm and its cationic charge density and hydrophobicity may be the key factors in transient pore formation. Repeatable poration could be obtained, and entry of  $\text{Ca}^{2+}$  via voltage-gated channels remains intact after application of low doses of poly-APS  
30 which indicates that at least at low concentrations cytotoxic damage does not occur as a result of pore formation.

Poly-APS produced highly variable  $\text{Ca}^{2+}$  transients, which

could reflect the nature of  $\text{Ca}^{2+}$  stores within DRG neurones and variable expression of voltage-activated  $\text{Ca}^{2+}$  channels. Poly-APS evoked membrane potential depolarisation would activate endogenous  $\text{Ca}^{2+}$  channels and both entry of  $\text{Ca}^{2+}$  through poly-APS pores and native channels would mobilise  $\text{Ca}^{2+}$  from stores through  $\text{Ca}^{2+}$ -induced  $\text{Ca}^{2+}$  release. Variability in responses to KCl-evoked depolarisation has been reported in DRG neurones but single neurones can respond consistently to repeated stimulation with KCl and poly-APS. Crude organic extracts containing bioactive pyridinium alkaloids from the sponge *Amphimedon viridis* show selective antibacterial activity. Some marine bacterial strains that possibly have symbiotic relations with the reef sponges were resistant to the toxin extracts. Poly-APS has not been found to have antibacterial activity against terrestrial and pathogenic gram-positive and gram-negative bacteria. Little comparative data has been obtained on the actions of sponge toxins on diverse pro- and eucaryotic cell types. However, our findings raise the possibility that variable sensitivity of cells and organisms to poly-APS is based on their intrinsic membrane composition and properties. The variability in the responses to poly-APS seen in the DRG neurone cultures could in part reflect the heterogeneous population of neurones in this preparation, with sensory neurones varying in their anatomical, biophysical and pharmacological characteristics.

It was also clearly demonstrated above that zinc, if continually present during application of poly-APS, could attenuate the effects of the sponge toxin preparation on membrane potential, input resistance, whole cell currents and  $\text{Ca}^{2+}$  permeability. This could involve an interaction between zinc and the cell membrane to prevent access of the toxin to sites on the membrane. The Shai-Matsuzaki-Huang model for

actions of antimicrobial peptides may provide a mechanism for pore formation by poly-APS. The initial stage in the model involves pore-forming molecules carpeting the outer membrane leaflet and this may be prevented by extracellular  $Zn^{2+}$  binding to negative charges on the membrane. The affinity of zinc could be much greater than that of pyridinium compounds and hence as seen in this study premixed poly-APS with zinc results in reduced poration. Thus zinc might prevent the further stages in pore formation, which are integration of the pore former into the membrane, thinning of the outer leaflet, phase transition and "wormhole" formation with the poly-APS molecules finally spanning the membrane producing a conductance pathway. Alternatively, zinc may interact with poly-APS directly to disrupt its biological activity. Previous work on erythrocytes has suggested that  $Zn^{2+}$  and other divalent cations ( $Hg^{2+}$ ) close pores produced by poly-APS but not lysis caused by hypoosmotic shock. This has not been demonstrated in this study, although, as suggested above, the interactions of poly-APS with membranes may vary and thus influence the protective actions of zinc, and the mechanism of erythrocyte lysis is made distinct by the osmotic influence of haemoglobin.

Fura-2 was used to measure changes in intracellular  $Ca^{2+}$  produced following pore formation by poly-APS, and significant reductions in fluorescence ratios were seen when zinc was applied. However, several features of the  $Ca^{2+}$  imaging experiments require further consideration. Firstly, fura-2 has also proved useful in measuring small increases in intracellular zinc because of its high zinc sensitivity ( $K_D$  values  $\sim 2$  nM), which is about 100 times lower than that for  $Ca^{2+}$ . The small sustained responses consistently observed with



zinc present seem to reflect fura-2 detecting zinc rather than  $\text{Ca}^{2+}$ , the sustained nature of these events indicates that unlike  $\text{Ca}^{2+}$ , zinc was not readily removed from the intracellular compartment. This suggests that zinc can pass  
5 through the poly-APS-evoked pores but that  $\text{Ca}^{2+}$  entry was attenuated to a greater extent than the data indicates. Combined with the electrophysiological data, the imaging studies with fura-2 indicate that zinc reduced levels of poration, and may also inhibit  $\text{Ca}^{2+}$  entry through the pores  
10 that are formed by poly-APS. However, it is possible that zinc interactions with fura-2 compete significantly to reduce  $\text{Ca}^{2+}$  detection. Secondly, zinc has been shown to inhibit voltage-activated neuronal conductances, including those carried by  $\text{Ca}^{2+}$  which may contribute to the poly-APS responses, and this  
15 could result in an overestimation of the inhibitory action of zinc on poly-APS effects. However, data has also been obtained from HEK 293 cells and under voltage clamp that are consistent with zinc inhibiting poly-APS-evoked pore formation independently of voltage-activated  $\text{Ca}^{2+}$  channels.

20

Poly-APS can reversibly form pores in the membranes of DRG neurones and HEK 293 cells, and applying extracellular zinc with the preparation of sponge toxins can attenuate these actions. The survival of cells even after dramatic but  
25 temporary changes in membrane potential and input conductances suggests that transient pore formation by low doses of poly-APS can be used to deliver materials to the intracellular environment, without cell damage.

30 The transient, reversible nature of the pores formed by the sponge toxins prompted further study to investigate whether these transient membrane pores/lesions were sufficient to conduct plasmid cDNA across the lipid bilayer.

In this connection, qualitative assessment of HEK293 cells incubated with 0.5 µg/ml poly-APS or halitoxin and 2.5 µg pEGFP indicated successful transfection, as a number of 5 fluorescent cells were observed (Fig. 11A, lower panels). This was in contrast to cells incubated with toxin alone (Fig. 11A upper panels) or pEGFP alone, which were not transfected and displayed no fluorescent cells. Use of a commercial transfection reagent (lipofectamine) to transfect HEK293 cells 10 (Fig. 11A, left-hand panels) provided a positive control for both the pEGFP cDNA and the imaging of transfected fluorescent cells. Extensive optimisation experiments were required to maximise transfection of HEK293 cells using sponge toxins and to provide a standardised protocol. Such experiments 15 investigated the influence on transfection efficiency of a number of variable factors and assessed both the percentage of cells transfected and cell survival.

A range of cDNA concentration experiments indicated that 2.5 20 µg pEGFP maximises transfection efficiency without inducing intolerable toxicity. The use of lesser quantities of cDNA, whilst being far less toxic, displayed low transfection efficiency, with greater quantities producing the opposite effect (high toxicity, moderate transfection efficiency).

25

Experiments focussing on the duration of incubation with toxin/cDNA revealed that a 5 min incubation with toxin prior to cDNA addition produced higher levels of transfection compared to simultaneous addition or longer periods of toxin 30 pre-incubation. Furthermore, a period of 180 min incubation was optimal following addition of cDNA, with shorter periods reducing transfection efficiency and longer periods proving too toxic. Serum has a profound effect on lipid-mediated cDNA

transfection, and so it was investigated whether similar effects would be seen with toxin-mediated transfection. Serum-free conditions proved most effective for toxin-mediated transfection, without raising toxicity levels, and in addition  
5 a return to standard 10% serum-containing medium following transfection was more beneficial than raising serum levels to 20% or above. Quantification of pEGFP transfection indicated that 0.5 µg/ml poly-APS/halitoxin achieved higher levels of transfection than either 0.05 µg/ml or 5.0 µg/ml (Fig. 11B).  
10 Comparable levels of transfection were achieved with both toxins, although at 0.5 µg/ml, poly-APS achieves marginally higher transfection success (Fig. 11B). These concentration-driven toxin activities coincide with the electrophysiological toxin profiles presented in Figures 4 and 5. Although these  
15 transfection levels are 4-fold lower than lipofectamine transfection efficiency, they are raised considerably above cDNA-only control levels (consistently 0). Analysis of the lethality of these toxins suggests that at both 0.05 µg/ml and 0.5 µg/ml neither of the toxins induce cell death above the  
20 tolerable level of death seen in lipofectamine-mediated transfection (typically 88-95% survival of HEK293 cells) (Fig. 11C). However, approximately 10 - 12 fold more death occurred with 5.0 µg/ml poly-APS and halitoxin (12% and 6% survival respectively) (Fig. 11C), suggesting a dose-dependent sponge  
25 toxin toxicity, with cells tolerant up to 0.5 µg/ml toxin. Furthermore, this very high toxicity explains the low proportion of fluorescent cells (Fig. 11B), with the toxin killing the cells as opposed to transfecting them at 5.0 µg/ml. This compliments data presented in Figures 5A and 5B  
30 concerning dose-dependent reduction in pore reversibility, which may be a factor contributing heavily to toxicity.

Stable transfection of HEK293 cells using 0.5 µg/ml poly-

APS further illustrates this toxin as a potentially useful transfection tool. Following selection of stable colonies in hygromycin B selection reagent, a large number of colonies remained in 6-well plates transfected using lipofectamine or 5 poly-APS, in contrast to control cells which had no hygromycin B resistant colonies (Table 2), indicative of hygromycin B lethality to normal cells.

Table 2. Comparison of lipofectamine and poly-APS transfected 10 stable colony number. Colonies of hygromycin B selected HEK293 cells stably transfected with pEGFP or TNFR2 cDNAs using poly-APS or lipofectamine were counted in culture wells prior to harvesting. Upper panel shows comparison of lipofectamine and poly-APS colony numbers, n = 5 and n = 10 respectively. Lower 15 panel split poly-APS stable data into either pEGFP or TNFR2 expressing colonies, n = 5 in both cases.

	Vehicle	Mean colony Number
	None	0
20	Lipofectamine	18.5 ± 4.5
	poly-APS	7.1 ± 1.4
	Poly-APS stable clones cDNA	
	pEGFP	6.2 ± 2.0
25	TNFR2	8.0 ± 2.0

In addition, although lipofectamine is more effective than poly-APS at stably transfecting HEK293 cells (2.5-fold more colonies in lipofectamine plates), a lower yet still relevant 30 level of transfection is achieved using poly-APS than commercial transfection reagents (Table 2 and Fig. 11B). Moreover, confocal visualisation of pEGFP stable clones

suggests larger numbers of positive cells in lipofectamine wells, compared to poly-APS transfected wells, although again poly-APS shows considerably larger numbers of stable cells compared to controls (Fig. 12A). The type of cDNA being co-  
5 introduced to the cell along with pBABE has no bearing on the success of stable expression, with poly-APS-mediated stable transfection of pEGFP and TNFR2 cDNA producing comparable numbers of stable colonies. Analysis and comparison of TNF-receptor levels in HEK293 cells, and two TNFR2 stably  
10 transfected clones, show relatively low and comparable labelling of TNFR1 in all three cell types (Fig. 12B), consistent with wild-type TNFR1 expression. In contrast, only clone 4 shows raised TNFR2 labelling, indicating successful stable transfection of TNFR2 cDNA. Clone 7 does not show an  
15 enhanced TNFR2 expression level, with very low labelling similar to control cells (Fig. 12B), suggesting only pBABE and not TNFR2 cDNA has been successfully transfected, resulting in expression of only the hygromycin resistant phenotype. Study of the same cells using FACS shows a similar rightward  
20 shift in fluorescence intensity for all three cell types when labelled with TNFR1-specific antibodies compared to secondary antibody alone (Fig. 12C), indicating similar levels of TNFR1 in the three cell types. However no TNFR2 shift is visible in untransfected or clone 7 cells with TNFR2 histogram lines  
25 overlying secondary antibody alone lines (Fig. 12C), suggesting little or no TNFR2 expression in the Hygromycin-resistance-only stable clone 7, as is seen in wild-type HEK293 cells. In higher expressing stable clone 4, a considerable increase in fluorescence intensity is apparent in TNFR2  
30 labelling, consistent with enhanced TNFR2 expression (Fig. 12C). Thus, clone 4 stably co-expresses both hygromycin resistant and TNFR2 enhanced phenotypes, following initial transfection with poly-APS, as proved by both confocal and

FACS analysis, demonstrating the usefulness of these reversible toxins as transfection reagents. There have been reports of the ability of keratinocytes to incorporate plasmid DNA without the use of transfection reagents. However, this natural DNA uptake process seems to be particular to the keratinocyte cell phenotype with surrounding fibroblasts unable to naturally incorporate foreign genetic information. The existence of plasmid DNA-binding proteins specific to keratinocytes was purported; however, it is clear that such mechanisms do not exist in HEK293 cells, as cDNA incorporation was not detected in the absence of transfecting reagent (Fig. 11A and 12A and Table 2).

The present invention provides that a simple incubation protocol using poly-APS can enable the introduction of large pieces of super-coiled DNA into living cells without irreparable cell damage. This novel and reproducible approach allows both transient and stable transfection conveniently with plasmid DNA, which has never before been achieved with pore forming molecules of this nature.

Consistent with findings in cultured dorsal root ganglion neurones and F-11 cells, halitoxin and poly-APS preparations produced conductance increases and raised intracellular  $\text{Ca}^{2+}$  in a manner consistent with pore or lesion formation in the cell membranes of HEK293 cells. It seems likely that the degree of polymerisation influences how readily the responses reverse; however, both sponge toxin preparations (halitoxin and poly-APS) contain two or more distinct alkyl pyridinium compounds. In addition to size of alkyl pyridinium compounds, issues of aggregation and lipid membrane constituents are likely influences on pore formation and cell recovery, and even between cells from a cell line considerable variability

in responses was observed.

The exact pore size formed by poly-APS and halitoxin preparations to allow transport of plasmid cDNA inside cells is not presently known, although it has been predicted to be 2.9 nm. It is known that the pore size must be greater than 2.6 nm as the maximal *Staphylococcus aureus*  $\alpha$ -toxin pore size is only sufficient to allow flux of ions and short (<200 bp) single-stranded oligonucleotides but not other biomolecules such as double-stranded DNA or protein. Moreover, the ionic components of *Staphylococcus aureus*  $\alpha$ -toxin hinder DNA movement through the pore of its multiple subunit form. Although only marginally larger in diameter, the estimated poly-APS aggregate pore does not contain a highly charged regions that would repel DNA and preclude its ability to allow plasmid DNA transfer into cells. Clearly the pore size must be greater than the width of a double-stranded  $\alpha$ -helical DNA chain (2 nm). The average predicted poly-APS pore size is 2.9 nm, and there may exist larger aggregated pores large enough to allow passage of soluble plasmid cDNA into cells. It could also be that cDNA in solution is capable of intertwining its  $\alpha$ -helices to minimise the overall diameter of the two double-stranded DNA chains that are side-by-side. In addition to the size of the pore formed, the transient nature of the pores that have been formed using poly-APS and halitoxin, are likely to be the reason for these reagent's successful use as transfection chemicals. Other pore-forming chemicals such as saponin and  $\alpha$ -escin produce pores in plasma membranes large enough to allow passage of sizeable 135 - 150 kDa proteins. These non-reversible pore-forming chemicals will be adequate for short-term experimentation, but cells exposed to them will eventually die through massive necrosis and  $\text{Ca}^{2+}$  excitotoxicity. The effective non-reversible nature of such

detergents and toxins is what renders them ineffective for DNA transfer in systems where the subsequent prolonged survival of the cell is necessary. The transfer of cDNA into HEK293 cells and subsequent protein expression indicated that both  
5 sponge toxin preparations at 0.5 µg/ml were able to provide reversible pores for cDNA transfer without cytotoxicity. The optimisation protocol indicated that pre-incubation with the sponge toxins and pore formation prior to addition of cDNA was useful, suggesting that in solution toxin and cDNA  
10 interactions attenuated pore formation and/or cDNA transfer. Similar effects were seen with serum present, which also reduced transfection efficiency achieved with poly-APS. Poly-APS has strong interactions with serum proteins and consistent with this study, incubation of poly-APS with highly diluted  
15 serum can prevent the hemolytic activity of poly-APS. Comparison between the methods using lipofectamine and sponge toxin preparations showed that although lipofectamine has higher efficiency, poly-APS has some clear advantages including high stability and good water solubility, and the  
20 sponge toxins provide a simpler and more consistent method which are of benefit to *in vivo* studies.

In summary, the above results show that viable cells transfected with DNA can be obtained using transient pore  
25 forming alkyl pyridinium sponge toxins and a simple pre-incubation protocol.

Poly-APS has also been used to deliver the cytoskeletal protein tau *in vitro* into cultures of hippocampal neurones and  
30 *in vivo* into the rat hippocampus. Hyperphosphorylated tau is a key characteristic of brains from patients with Alzheimer's disease and the possibility of generating models of tauopathies has been investigated. Tau protein induced



deficits were evaluated using electrophysiology,  $\text{Ca}^{2+}$  imaging and water maze learning. The intracellular loading of preparations with the protein was assessed using tau specific antibodies. Cultured hippocampal neurones loaded with tau  
5 survived for at least 24 hours and normal action potentials and  $\text{Ca}^{2+}$  transients could be recorded from them. Similarly *in vivo*, animals performed water maze learning and memory tests normally after delivery of tau to the hippocampus. However, if in addition to delivering tau, cultures were treated with  
10 the phosphatase inhibitor okadaic acid, action potentials and clear  $\text{Ca}^{2+}$  transients could not be evoked in the few neurones that survived. Furthermore, animals treated *in vivo* with poly-APS, tau and okadaic acid could learn the water maze platform position, but had impaired reversal learning when the platform  
15 position was changed. Long-term potentiation measured in hippocampal slices *ex vivo* was also attenuated.

Site specific delivery of tau to intracellular compartments followed by hyperphosphorylation provides a new approach to the investigation of cellular aspects of  
20 Alzheimer's disease. These data also provide further proof that poly-APS can be used to deliver macromolecules into cells to achieve transfection or model disease states *in vitro* and *in vivo*.

25 Water maze experiments involved training animals to find a platform in a quadrant of a circular bath. Once the animal had learnt the position of the platform the platform could be moved to test the animals ability to show reversal learning.

30 Figure 13 shows two trails and the swimming paths taken to the platform. In trail 1 the animal swims round the bath and has not learnt where the platform is but during trial 16 the animal swims directly to the platform.

Poly-APS (sponge toxin) was applied to the hippocampus either alone or with tau, tau was also applied without poly-APS. A mini pump was then used in some animals to deliver okadaic acid over 7 days, this blocks phosphatases resulting in 5 hyperphosphorylation of proteins including tau protein.

In figure 14 the data for learning where the platform is for each group of experimental animals is shown. All animals learnt where the platform was, as reflected by the time (above 10 25%) spent in the quadrant with the platform and the swimming path lengths. When the position of the platform was moved, the animals that had been treated with poly-APS, tau and okadaic acid failed to show learning reversal (figure 15). All other treatment groups did.

15

As can be seen from Figure 16, which shows ex-vivo data, hippocampal slices were used to study long term potentiation (LTP). LTP is a form of neuronal plasticity which last many hours or days. If pathways get tetanic stimulation (arrow) 20 subsequent electrophysiological signalling is increased. Brains from animals treated with poly-APS, Tau and okadaic acid showed less LTP.

As can be seen from Figure 17, tau was successfully delivered 25 to the intracellular compartments of hippocampal neurones in culture. The tau level (black) were detected using specific antibodies to this protein.

As shown in Figure 18, cultured hippocampal neurones treated 30 with poly-APS and tau produced action potentials, whilst neurones treated with poly-APS, tau and okadaic acid failed to show action potential firing even when stimulated with much larger currents.

As shown in Figure 19, Cultured hippocampal neurones were stimulated with KCl pulses. Neurones treated with okadaic acid as well as poly-APS and tau failed to show proper calcium transients. With the data shown in Figure 18, this shows that  
5 the excitability of these neurones was inhibited probably by misfunctioning of voltage-activated channels.

As shown in Figure 20, the mean data for calcium transient for neurones exposed to the different treatments, clearly  
10 distinguished the neurones exposed to poly-APS, tau and okadaic acid from the neurones exposed to poly-APS and okadaic acid, and poly-APS and tau. With the data from Figures 18 and 19, this shows that the excitability of these neurones was greatly reduced and probably involved  
15 misfunctioning of voltage-activated channels.

Claims

1. Use of a composition comprising a sponge toxin for the reversible formation of a membrane pore.

5

2. Use according to claim 1, wherein the sponge toxin comprises at least one polymeric 1,3-alkylpyridinium salt (poly-APS).

10

3. Use according to either claim 1 or claim 2 wherein the sponge toxin is obtained from the sponge *Reniera sarai*, *Callyspongia ridleyi*, *Haliclona erina*, *Haliclona rubens*, *Haliclona viridis*, *Amphimedon viridis*, *Callyspongia fibrosa* and *Amphimedon compressa*.

15

4. Use according to any preceding claim, wherein the sponge toxin has a molecular weight of between 5 kDa and 20 kDa.

20

5. Use according to claim 4, wherein the sponge toxin has a molecular weight of 5.5 kDa or 18.9 kDa.

6. Use according to any preceding claim, wherein the concentration of sponge toxin is between 0.5 ng/ml and 5.0  
25 µg/ml.

7. Use according to claim 6 wherein the concentration of sponge toxin is between 0.5 ng/ml and 0.5 µg/ml.

30

8. A method for the reversible formation of membrane pores, the method comprising the steps of:-

a) incubating the membrane in the presence of a composition according to any preceding claim; and

b) removing the composition from contact with the membrane.

9. A method according to claims 8, wherein zinc solution is added to attenuate the formation of the membrane pore.

10. A method according to claim 9 wherein the concentration of zinc solution is between substantially 1 to 10 2 mM.

11. A method according to claim 9 or 10, wherein the concentration of zinc is 1.5mM.

15 12. A method for transfection of a macromolecule into a cell *in vitro*, the method comprising the steps of:-  
a) incubating the cell in the presence of a composition comprising a sponge toxin;  
b) removing the composition from contact with the membrane;  
20 and  
c) adding the macromolecule.

13. A method according to claim 12, wherein the macromolecule is cDNA, protein, peptide, lipid or 25 oligonucleotide.

14. A method according to claim 12 or 13, wherein the cell is incubated in the presence of the composition for between 1 and 20 minutes prior to addition of the 30 macromolecule.

15. A method according to any one of claim 12 to 14 wherein the cell is incubated in the presence of the

composition for 5 minutes prior to the addition of the macromolecule.

16. A method according to any of claims 13 to 15, wherein between 1.0 and 5.0 µg nucleic acid is added.

17. A method according to any of claims 13 to 16, wherein 2.5 µg nucleic acid is added.

10 18. A method according to any of claims 12 to 17, comprising incubating the cell, in the presence of the composition and macromolecule and replacing the composition and macromolecule with standard media.

15 19. A method according to claim 18 wherein the cells are incubated for between 20 and 200 minutes.

20 20. A method according to either claim 18 or 19 wherein the cells are incubated for 180 minutes.

21. A method for transfection of a macromolecule into a cell *in vivo*, the method comprising the step of:-

a) incubating the cell in the presence of a composition comprising a sponge toxin and the macromolecule.

25

22. A method according to claim 21, wherein the macromolecule is cDNA, protein, peptide, lipid or oligonucleotide.

30 23. A method according to claim 21 or 22, wherein the macromolecule is the cytoskeletal protein tau.

24. A method according to any of claims 21 to 23 wherein

the cell is a hippocampal neurone.

25. A model for use in the study of neurological disease or treatments thereof, the model comprising a rodent having  
5 undergone application of a composition comprising a sponge toxin, tau protein and phosphatase inhibitor to the hippocampus.

26. A model according to claim 25 wherein the  
10 neurological disease is Alzheimer's.

27. A model according to claims 25 or 26 wherein the rodent is a rat or a mouse.

15 28. A method of studying a neurological disease, the method comprising:

a) applying a composition comprising a sponge toxin, tau protein and phosphatase inhibitor to the hippocampus of a rodent; and

20 b) studying the effect on the rodent.

29. A model according to claims 25 to 27 or a method according to claim 28 wherein the phosphatase inhibitor is okadaic acid.




Fig 1A




Fig 1B




Fig 1C






Fig 2A




Fig 2B

3/20




Fig 3A




Fig 3B




Fig 3C

4/20




Fig 4A




Fig 4B




Fig 4C








Fig 4D

5/20



6/20




Fig 6A




Fig 6B






Fig 6C

7/20



8/20




Fig 8A




Fig 8B




Fig 8C

9/20




Fig 9A




Fig 9B



10/20

Fig 10A





Fig 10B



11/20

Fig 11A




Fig 11B





Fig 11C



12/20

Fig 12A




Fig 12B









Fig 12C



# Water Maze Experiments- Learning

13 / 20

Fig. 13



Trial 1

Trial 16

# Water Maze - All groups learn the platform position




Fig. 14B All groups display spatial bias




Fig. 14C




Fig. 14C

All groups learn the platform position

Water Maze - In reversal learning tests the group treated with poly-APS +  $\tau$  + okadaic acid show no spatial bias

Fig. 15 A






Fig. 15 B



Treatment with poly-APS +  $\tau$  + Okadaic acid reduced  
Long Term Potentiation


Fig. 16



Poly-APS can be used to load  $\tau$  into hippocampal neurones in culture

Fig. 17

Poly-APS +  $\tau$




Control



# Hyperphosphorylated $\tau$ disrupts action potentials in cultured hippocampal neurones


Fig. 18

Poly-APS +  $\tau$       Poly-APS +  $\tau$  + Okadaic acid




# Hyperphosphorylated $\tau$ disrupts electrophysiological properties of hippocampal neurones in culture

Fig. 19




Hyperphosphorylated tau disrupts electrophysiological properties of hippocampal  
neurons in culture

Fig. 20



**PCT/GB2004/002659**



**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

☒ **BLACK BORDERS**

☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**

☐ **FADED TEXT OR DRAWING**

☒ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**

☐ **SKEWED/SLANTED IMAGES**

☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**

☐ **GRAY SCALE DOCUMENTS**

☒ **LINES OR MARKS ON ORIGINAL DOCUMENT**

☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**

☐ **OTHER:** \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**